








ГЕНЕТИКА GENETICS



DOI: 10.18413/2658-6533-2024-10-2-0-1

УДК 575.174.015.3:616.831-005.4-056.83

HSPD1 gene polymorphism is associated with an increased risk of ischemic stroke in smokers

Tatiana A. Stetskaya¹ , Ksenia A. Kobzeva¹ , Sergey M. Zaytsev² ,
Irina V. Shilenok³ , Galina V. Komkova¹ , Natalia V. Goryainova¹ ,
Olga Yu. Bushueva¹ 

¹ Kursk State Medical University,

3 Karl Marx St., Kursk, 305004, Russia

² Pavlov First Saint Petersburg State Medical University,

6-8 Leo Tolstoy St., Saint Petersburg, 197022, Russia

³ Kursk City Clinical Emergency Hospital,

14 Pirogov St., Kursk, 305035, Russia

Corresponding author: Olga Y. Bushueva (olga.bushueva@inbox.ru)

Abstract

Background: Ischemic stroke (IS) is a leading cause of death and disability worldwide. Since the influence of vascular stressors prompts cross-reactivity with microbial and human HSPD1 (HSP60), which results in the degradation of vascular endothelium and intimal infiltration by mononuclear cells, HSP60 is viewed as a key player in the autoimmune theory of atherosclerosis. **The aim of the study:** The present pilot study investigated whether the tag SNP rs11682567 *HSPD1* is associated with the risk of IS. **Materials and methods:** DNA samples from 1176 unrelated Russian subjects (708 IS patients and 468 healthy controls) were genotyped for tagging SNP rs11682567 in the *HSPD1* gene using the MassArray-4 system. A log-additive regression model was used to interpret the associations between genotypes and IS risk. **Results:** We discovered that smoking, a major environmental endothelial stressor, dramatically modifies the link between rs11682567 and IS, with a relationship only being seen in smokers (risk allele G; OR = 2.11, 95% CI = 1.17-3.82; P = 0.0075). The functional annotation of this SNP revealed that risk allele G of rs11682567 impacts the expression and alternative splicing of genes that regulate brain ischemia-reperfusion injury, reactive oxygen species production, toll-like receptor 4 internalization, and pro-inflammatory properties. Moreover, allele G rs11682567 creates binding sites for transcription factors jointly involved in biological processes encompass neurogenesis, monocyte differentiation, fatty acid metabolism, cell signalling, and apoptosis. **Conclusion:** Thus, rs11682567 SNP in the *HSPD1* gene represents a novel genetic marker of ischemic stroke, whose significance for the pathogenesis of IS is mediated by smoking.

Keywords: ischemic stroke; heat shock; mitochondrial folding; *HSPD1*; HSP60; chaperones; atherosclerosis; autoimmune response; rs11682567; smoking

For citation: Stetskaya TA, Kobzeva KA, Zaytsev SM, et al. *HSPD1* gene polymorphism is associated with an increased risk of ischemic stroke in smokers. *Research Results in Biomedicine*. 2024;10(2):175-186. DOI: 10.18413/2658-6533-2024-10-2-0-1

Introduction. Ischemic stroke is a major global cause of death and disability [1]. A lot of studies have been done so far to investigate the genetic nature of stroke [2] and stroke-related phenotypes, like hypertension [3] and atherosclerosis [4]. Heat Shock Proteins (HSPs) are engaged in a number of cellular processes, including immunological response, inflammation, and apoptosis, that may have an effect on the outcomes of stroke [5]. Also, by stabilising proteins and assisting in protein re-folding and degradation, HSPs directly contribute to the protection of cells and tissues against the damaging effects of ischemia [6].

HSPD1, otherwise known as heat shock protein 60 (HSP60), is a chaperonin that is present in human mitochondria. It assists in protein folding and is required for preserving mitochondrial functionality, which is essential for cell survival [7]. A major risk factor for stroke is atherosclerosis, an inflammatory condition which was initially referred to an immunological response against HSP60 [8]. There is a

chance of cross-reactivity between human and microbial HSP60, which can cause endothelial damage and inflammation and possibly aid in the onset of atherosclerosis [4]. To our knowledge, no research has looked into the relationship between *HSPD1* SNPs and IS risk.

The aim of the study. To investigate whether SNP rs11682567 *HSPD1* is related with the risk of IS.

Materials and methods. The study enrolled 1176 unrelated Russians from Central Russia, including 708 patients with IS and 468 healthy people. The study protocol was approved by the Kursk State Medical University's Ethical Review Committee. Before being accepted into the study, each participant provided written informed consent and agreed to the inclusion criteria listed below: self-declared Russian ancestry and birthplace in Central Russia.

In Table 1, the study population's baseline characteristics are presented.

Table 1

Baseline characteristics of the studied groups

Baseline characteristics		IS patients (n=708)	Controls (n=468)	P-value
Age, Me [Q1; Q3]		63 [55; 70]	57 [54; 64]	<0.001
Gender	Males, N(%)	399 (56.4%)	168 (35.9%)	<0.001
	Females, N(%)	309 (43.6%)	300 (64.1%)	
Smoking	Yes, N (%)	365 (51.5%)	82 (17.5%)	<0.001
	No, N (%)	343 (48.5%)	386 (82.5%)	

Both the Regional Vascular Centre of Kursk Regional Clinical Hospital and the Neurology Clinics of Kursk Emergency Medicine Hospital participated in the study's patient recruitment [9-12]. In order to diagnose IS during the acute phase of stroke, the results of the neurological examination, brain computed tomography, and/or magnetic resonance imaging were used. The following conditions were used to exclude individuals from the IS study: traumatic brain injury; intracerebral haemorrhage; hemodynamic or dissection-related stroke; hepatic or renal failure; autoimmune, oncological, or other

disorders that can produce an abrupt cerebrovascular event. All of the IS patients had a history of hypertension and were taking medication to lower their blood pressure.

The control group consisted of healthy volunteers with normal blood pressure, not receiving antihypertensive therapy, without clinical signs of cardiovascular, cerebrovascular and other serious diseases. If a healthy individual's systolic blood pressure was less than 130 mm Hg and/or their diastolic blood pressure was less than 85 mm Hg on at least three different occasions, they would have qualified for

inclusion in the control group. When undergoing medical examinations at public institutions and industrial businesses in the Kursk region, control subjects were recruited from Kursk hospitals during particular periods [13, 14]. During the same time frame and from the same population, this cohort was recruited [15].

The selection of SNPs was based on the following criteria: the SNP needs to be tagging, with a minor allele frequency of at least 0.05 in the European population, and stand out for having a high regulatory potential. The bioinformatic tools SNPinfo Web Server (<https://snpinfo.niehs.nih.gov/>) and LD TAG SNP Selection, which were used to identify SNPs based on the reference haplotypic structure of the Caucasian populations (CEU, TSI) of the project HapMap, determined that the rs11682567 *HSPD1* is a tagging SNP. This SNP is located in intron chr2:197490429 (GRCh38.p14) and characterized by the replacement T>G (<https://www.ncbi.nlm.nih.gov/snp/rs11682567>).

The regulatory potential of rs11682567 was evaluated using a variety of bioinformatic tools [16]. The RegulomeDB online tool (<https://regulomedb.org/regulome-search/> (accessed November 20, 2022)) showed that rs11682567 is characterized by a regulatory coefficient of 5 (TF binding or DNase peak). This SNP is characterized by Enhancer histone marks and regulatory motifs, according to the HaploReg (v4.1) database (<http://archive.broadinstitute.org/mammals/haploreg/haploreg.php> (accessed on November 20, 2022)).

The NCBI source's data (<https://www.ncbi.nlm.nih.gov/snp/>) indicates that this genetic variation is described by an average minor allele frequency of >0.05 in European populations. In light of the fact that SNP rs11682567 satisfies the prerequisites for study inclusion, it was chosen for our research.

Genetic Analysis. The Laboratory of Genomic Research at the Research Institute for Genetic and Molecular Epidemiology of Kursk State Medical University (Kursk, Russia) performed the DNA analysis. The conventional technique of phenol/chloroform extraction and

ethanol precipitation was used to extract genomic DNA from thawed blood samples. The MassARRAY-4 genetic analyzer (Agena Bioscience, San Diego, CA, USA) was used for genotyping.

The SNP primers were created through the use of the Agena Bioscience's Assay Design Suite software, and genotyping was accomplished by means of matrix-assisted laser desorption ionization mass spectrometry (MALDI-TOF). The remaining dNTP and primers were removed by employing the SAP enzyme after a PCR reaction specific to the locus. Mass-modified dideoxy-nucleotide terminators of an oligonucleotide primer were then utilized to carry out single-base extension, which occurred upstream of the desired polymorphic site.

A PCR reaction was conducted on a spectral chip using a 5 μ L reaction mixture that included 3 μ L of PCR mix consisting of 80.6 μ L of ddH₂O, 50.4 μ L of 10x buffer, 40.3 μ L of MgCl₂ (25 mmol/L), 10.1 μ L of a mixture of dNTP (25 mmol/L), 100.8 μ L of a mixture of PCR primers (0.5 mmol/L), and 20.16 μ L of HotStartTaq polymerase (5 U/ μ L), as well as 2 μ L of DNA (10 ng/ μ L). The reaction was carried out for 125 minutes in a CFX96 Bio-Rad thermal cycler, with the following program: denaturation at 95 °C for 2 minutes, followed by 44 cycles of denaturation at 95 °C for 30 seconds, annealing at 56 °C for 30 seconds, and elongation at 72 °C for 60 seconds. The samples were then incubated at 72 °C for 5 minutes.

The unreacted dNTPs present in the PCR samples were eliminated using the shrimp alkaline phosphatase (SAP) reaction, which involved the hydrolysis of dNTPs to nucleosides and phosphates by the alkaline phosphatase of shrimp. To achieve this, 2 μ L of SAP mixture, consisting of 161.6 μ L of ddH₂O, 18 μ L of 10xTS buffer, and 31.7 μ L of SAP enzyme, was added to each well, and the plate was loaded into the CFX96 Bio-Rad thermal cycler. The SAP reaction was carried out for 55 minutes, comprising 45 minutes at 37 °C, followed by 10 minutes at 85 °C.

To prepare for the third iPLEX reaction, an E-primer mix was created based on the molecular weight of the E-primers. The resulting mixture was evaluated using mass spectrometry to determine the intensity of the E-primer peaks, and the necessary quantities of E-primers were added to the mix to achieve the same signal intensity. 2 μ L of iPLEX-mix (consisting of 65.4 μ L of ddH₂O, 21.2 μ L of iPLEX buffer, 21.2 μ L of terminating mixture, 99.3 μ L of E-primer mix, and 4.33 μ L of iPLEX enzyme) was then added to each well. The plate was loaded into a CFX96 Bio-Rad thermal cycler and underwent denaturation at 94 °C for 30 seconds, followed by 40 cycles of denaturation at 94 °C for 5 seconds, annealing at 52 °C for 5 seconds, and elongation at 80 °C for 5 seconds. The reaction was then incubated at 72 °C for 3 minutes. After the iPLEX reaction was complete, 30 μ L of water was added to all wells, and the plate was loaded into a mass spectrometer.

To prevent the formation of contaminant peaks resulting from the binding of small polar molecules and positively charged sodium and potassium cations to oligonucleotides, the amplicons underwent robotic desalination using SpectroCLEAN resin (Agena Bioscience).

The samples were loaded onto a spectral chip with an ionization matrix using the Nano-dispenser workstation in automatic mode. The samples formed crystals on the SpectroCHIP, which were then ionized by a high-energy laser, resulting in the ionization of DNA molecules. Subsequently, the ionized DNA samples were passed through a time-of-flight tube with an electric field. By detecting the time-of-flight of analytes, we could determine the genotype of the analytes. Finally, the results of mass spectrometric analysis for each SNP were reviewed using the TyperAnalyzer program from Agena Bioscience.

To ensure quality control, 95 randomly selected DNA samples were chosen for repeat genotyping on the same platform, without knowledge of the case-control status. The repeatability test revealed a 100% concordance rate.

- QTLbase Bioinformatic tool (<http://www.mulinlab.org/qtlbase/index.html>

(accessed February 24, 2023)) was used for analysis of expression quantitative trait loci (eQTLs) in brain, whole blood, and blood vessels. Additionally, for QTL analysis of the peripheral blood effect the eQTL online tool was used.

- STRING database's bioinformatic tools were utilised to analyse the main functional partners of HSPD1 (<https://string-db.org/> (accessed February 24, 2023)). The STRING database was also used for analysis of biological processes and molecular functions reflecting interactions HSPD1 with key functionally relevant proteins (accessed February 21, 2023).

- The effect of rs11682567 *HSPD1* on the binding of transcription factors (TFs) to DNA based on the presence of the reference/alternative alleles was assessed using the atSNP Function Prediction online tool (<http://atsnp.biostat.wisc.edu/search> (accessed February 21, 2023)). In accordance with a positional weight matrix-based calculation of the impact of SNPs on how well TFs interact with DNA, certain TFs were added.

- It was possible to assess the joint participation of TFs linked to the reference/SNP alleles in overrepresented biological processes directly associated with the pathogenesis of IS using the Gene Ontology web tool (<http://geneontology.org> (accessed February 24, 2023)).

- The Comparative Toxicogenomics Database (CTD) resource (<http://ctdbase.org> (accessed February 24, 2023)) was used for interpretation on environment-associated correlates of *HSPD1*. Based on information from globally published studies, CTD enables the analysis of particular interactions between genes and chemicals in vertebrates and invertebrates. One chemical and one gene or protein were the subjects of this tool's analysis of binary interactions.

- For a bioinformatic examination of the connections between rs11682567 *HSPD1* and stroke-related phenotypes, intermediate phenotypes (such as blood pressure, heart rate etc.), the Cerebrovascular Disease Knowledge Portal (CDKP) (<https://cd.hugeamp.org/> (accessed

February 24, 2023)) and Cardiovascular Disease Knowledge Portal (<https://cvd.hugeamp.org/> (accessed February 24, 2023)) were used.

Results and discussion
Associations of rs11682567 HSPD1 with the Risk of Ischemic Stroke. Table 2 displays the genotype frequencies of rs11682567 *HSPD1* in the study groups.

Table 2

Results of the analysis of associations between rs11682567 HSPD1 gene SNP and ischemic stroke risk

Genotypes	Controls (N=468)	IS patients (N=708)	OR (95% CI)	P
Entire group				
T/T	360 (76.9%)	511 (72.2%)	adj1.25 (0.95-1.65)	adj0.11
G/T	104 (22.2%)	184 (26%)		
G/G	4 (0.8%)	13 (1.8%)		
Maf (G)	0.120	0.148		
Nonsmokers				
T/T	292 (75.7%)	256 (74.6%)	1.07 (0.78-1.46)	0.68
G/T	90 (23.3%)	82 (23.9%)		
G/G	4 (1%)	5 (1.5%)		
Maf (G)	0.127	0.134		
Smokers				
T/T	68 (82.9%)	255 (69.9%)	2.11 (1.17-3.82)	0.0075
G/T	14 (17.1%)	102 (27.9%)		
G/G	0 (0%)	8 (2.2%)		
Maf (G)	0.085	0.162		

Note: ¹ – odds ratio and 95% confidence interval; P – P-level. Every computation was done in relation to the minor alleles. Differences that are statistically significant are highlighted in bold. adj – calculations, performed with adjustment for sex, age, smoking.

In each case and control groups, the genotype frequencies of all examined SNPs were distributed according to the Hardy-Weinberg equilibrium ($P > 0.05$). The analysis of the total sample did not reveal associations between rs11682567 *HSPD1* and IS risk (Table 2).

Subsequent sub-group analysis showed that rs11682567 was associated with an increased risk of IS exclusively in smokers (risk allele G; OR = 2.11; 95% CI = 1.17–3.82; $P = 0.0075$).

Bioinformatics Analysis. The *HSPD1* gene is highly expressed in brain tissues, blood vessels, and whole blood. In brain tissues, gene expression levels of *HSPD1* (MeTPM) varies from 77.57 to 159.3; in blood vessels – from 122.4 to 131.0; in whole blood MeTPM=30.69.

QTL-effects. According to the QTLbase, the G allele rs11682567 is associated with

eQTL-mediated up-regulation of *HSPD1* expression in the brain-hippocampus ($\beta=0.174$; FDR=0.002), as well as up-regulation of *ANKRD44* in Blood-T cell CD4+naive ($\beta=0.02$; FDR=0.03) and *ANKRD44-IT1* in blood-T cells CD4+ naive ($\beta=0.038$; FDR=0.02)/blood-neutrophils CD16+ ($\beta=0.015$; FDR=0.036) (<http://www.mulin-lab.org/qtlbase>).

Furthermore, using the eQTLGen resource, it was discovered that the G allele rs11682567 had an impact on the expression of the genes *AC013264.2*, *MARS2*, *RFTN2*, *COQ10B*, *SF3B1*, and *HSPE1* as well as an increase in the expression of the *PLCL1* gene in peripheral blood (Table 3).

Using the bioinformatics tool FIVEx, it was feasible to determine how the G rs11682567 allele affects the alternative splicing of the *HSPD1*, *ANKRD44*, *COQ10B*,

HSPE1, *MOB4*, *RFTN2*, and *SF3B1* genes in blood cells and brain tissues by cis-sQTL-mediated effects (<https://fivex.sph.umich.edu>).

Analysis of transcription factors. Risk allele G of rs11682567 creates binding sites for transcription factors BRCA1, COMP1, FOXO3, FOXO4, HMGA1, IRF7, MYC, RUNX2, SOX10, SOX17, SOX2, Sox6, SOX9, SRY. According to the Gene Ontology resource, these transcription factors are jointly involved in overrepresented biological processes, such as «glial cell fate commitment» (GO:0021781; FDR=0.005); «regulation of monocyte differentiation» (GO:0045655; FDR=0.01); «neuronal stem cell population maintenance» (GO:0097150; FDR=0.014); «negative regulation of fatty acid metabolic process» (GO:0045922; FDR=0.03); «negative regulation of neuron differentiation»

(GO:0045665; FDR=0.003); «oligodendrocyte differentiation» (GO:0048709; FDR=0.003); «cellular response to growth factor stimulus» (GO:0071363; FDR=0.02); «cellular response to cytokine stimulus» (GO:0071345; FDR=0.009); «negative regulation of apoptotic process» (GO:0043066; FDR=0.02). The etiopathogenesis of IS is greatly influenced by all of these biological processes. The fact that TFs that bind to the G allele also control «DNA methylation» (GO:0044030; FDR=0.016) and «chromatin organization» (GO:0006325; FDR=0.004) adds to the notion that rs11682567 *HSPD1* plays a substantial role in the mechanisms of epigenetic regulation. It is remarkable that no overrepresented GO was discovered when TFs that bind to the reference allele T rs11682567 were examined.

Table 3

Cis-eQTL-mediated effects of rs11682567 *HSPD1* in peripheral blood according to the resource eQTLGen (<https://www.eqtngen.org/>)

SNP	Effect allele	Symbol	P-value	FDR	Z-score	N samples
rs11682567	G	<i>PLCL1</i>	4.44×10^{-149}	<0.0001	26.0043	31644
rs11682567	G	<i>AC013264.2</i>	1.24×10^{-71}	<0.0001	-17.8972	5502
rs11682567	G	<i>MARS2</i>	1.86×10^{-41}	<0.0001	-13.4873	31644
rs11682567	G	<i>RFTN2</i>	1.04×10^{-36}	<0.0001	-12.6552	31176
rs11682567	G	<i>COQ10B</i>	2.67×10^{-17}	<0.0001	-8.4598	31684
rs11682567	G	<i>SF3B1</i>	2.90×10^{-11}	<0.0001	-6.6516	31684
rs11682567	G	<i>HSPE1</i>	2.24×10^{-8}	0.0001	-5.5922	14263

Bioinformatic analysis of the associations of rs11682567 *HSPD1* with cerebral stroke and intermediate phenotypes. According to the bioinformatic resources Cerebrovascular Disease Knowledge Portal (CDKP) and Cardiovascular Disease Knowledge Portal (CVDKP), which integrate and evaluate the genetic association data from the greatest consortiums for the research of cardiovascular and cerebrovascular disorders, the risk allele G of rs11682567 *HSPD1* is associated with increasing in diastolic blood pressure (beta=0.007; P=0.0007), total cholesterol (beta=0.0033; P=0.033), coronary artery disease in subjects without diabetes (OR=1.09; P=0.043).

Protein-Protein Interactions. Using the STRING database, 10 proteins (proteins of the first shell of interactors) distinguished by the most pronounced interactions with *HSPD1* were discovered: *HSPE1*, *HSPA9*, *TLR4*, *TLR2*, *HSP90AA1*, *HSP90AB1*, *TST*, *GRPEL1*, *TRAP1*, *DHFR* (PPI enrichment p-value= 7.46×10^{-5}) (Figure).

It is interesting that the majority of *HSPD1*'s primary functional partners exhibit chaperone and co-chaperone activities, which are strongly associated with proteostasis (by regulation of protein import, misfolding and refolding of unfolded proteins under stress), inflammatory response (by NF-kappa-B activation), folate metabolism (Table 4).




Fig. Predicted Functional Partners of *HSPD1* (<https://string-db.org/>)

HSPD1 and its primary functional partners jointly participate in biological processes which reflects biological processes crucial to the pathophysiology of IS, such as proteostasis, oxidative stress, inflammation, regulation of extracellular matrix, endothelial dysfunction, heat stress, apoptosis, neurogenesis, cell signaling (<https://string-db.org/cgi/network?taskId=bACsG8aUeF2I&sessionId=bjoEZTIHrwUn>).

The current study demonstrates for the first time that polymorphism rs11682567 in the gene encoding Heat Shock Protein Family D (Hsp60) Member 1 (*HSPD1*) is linked to the risk IS and is significantly modified by smoking. *HSPD1* is expressed in vessels, brain tissues and blood. This chaperonin is responsible for regulating mitochondrial protein homeostasis and mitochondrial function: it encourages unfolded polypeptides to fold normally in stressful conditions, avoids misfolding, and facilitates imported proteins fold properly [17]. Previous studies have demonstrated a substantial function for *HSPD1* in the emergence of

cardiovascular diseases. Patients with CVD showed elevated HSP60 levels [18]. Currently, HSP60 is seen as a major player in the autoimmune theory of atherosclerosis [8, 19]. The highly conserved microbial HSP60 antigen is a target of protective, advantageous adaptive immunity that all humans develop [20]. However, exposure to endothelial stressors, such as components of cigarette smoke, causes *HSP60* expression in vascular endothelial cells that do not typically express *HSP60*, which causes a (cross)reaction and destruction of these target cells by preexisting cellular and humoral immunity against HSP60, resulting in intimal infiltration by mononuclear cells [19]. HSP60 also has a sizable impact on the regulation of vascular smooth muscle cells because it activates the IKK/NF- κ B, ERK/MAPK, and TLR4 pathways, which promote vascular smooth muscle cell migration, thicken the intima, and are important in vascular wall remodeling [18, 21, 22, 23]. Thus, in total, exposure to vascular stressors triggers a "vicious circle" of an autoimmune response leading to the progression of atherosclerosis.

Table 4

Main functional characteristics of HSPD1 predicted functional partners (<https://string-db.org/>)

Protein	Functions	*						Score
		1	2	3	4	5	6	
HSPE1	10 kDa heat shock protein, mitochondrial; Co-chaperonin implicated in mitochondrial protein import and macromolecular assembly. May also prevent misfolding and promote the refolding of unfolded polypeptides generated under stress conditions in the mitochondrial matrix.	•	•	•	•		•	0.999
HSPA9	Stress-70 protein, mitochondrial; Chaperone protein which plays an important role in mitochondrial iron-sulfur cluster biogenesis. May be involved in the regulation of proliferation of cells and cellular ageing.	•		•	•	•	•	0.999
TLR4	Toll-like receptor 4; operates through MYD88, TIRAP, and TRAF6, triggering the release of cytokines and the inflammatory response. In complex with TLR6, promotes sterile inflammation in monocytes/macrophages in response to oxidized low-density lipoprotein.				•	•	•	0.999
TLR2	Toll-like receptor 2; activates NF-kappa-B, causes the release of cytokines, and triggers the inflammatory response by acting through MYD88 and TRAF6. In response to the lipid moiety of lipoproteins, may also cause immune cells to become active and promote apoptosis.				•	•	•	0.998
HSP90AA1	Molecular chaperone that engages with a range of client protein classes via its interaction with various co-chaperone proteins or complexes, that act as adapters, simultaneously able to interact with the specific client and the central chaperone itself. Plays a critical role in mitochondrial import.			•	•		•	0.998
HSP90AB1	Molecular chaperone that promotes the maturation, structural maintenance and proper regulation of specific target proteins involved for instance in cell cycle control and signal transduction.			•	•		•	0.990
TST	Together with MRPL18, acts as a mitochondrial import factor for the cytosolic 5S rRNA. Only the nascent unfolded cytoplasmic form is able to bind to the 5S rRNA.				•		•	0.989
GRPEL1	GrpE protein homolog 1, mitochondrial; Seems to control the nucleotide-dependent binding of mitochondrial HSP70 to substrate proteins.	•		•	•	•	•	0.987
TRAP1	A negative regulator of mitochondrial respiration able to modulate the balance between oxidative phosphorylation and aerobic glycolysis.			•	•		•	0.986
DHFR	Dihydrofolate reductase; Key enzyme in folate metabolism.	•		•	•		•	0.984

Note: Data obtained with STRING database; * - Proven interaction mechanisms: 1 – Neighborhood; 2 – Cooccurrence; 3 – Coexpression; 4 – Experiments; 5 – Databases; 6 – Textmining.

According to bioinformatics analysis utilising the STRING database, the key functional partners of HSPD1 are involved in a number of overrepresented biological processes associated with atherosclerosis, managing not just proteostasis but also oxidative stress, MyD88-dependent toll-like receptor signaling pathway, I-kappaB phosphorylation, production of pro-inflammatory cytokines, nitric-oxide synthase level, macrophage activation and proliferation, extracellular matrix organization, apoptosis, tumor necrosis factor production, neurogenesis. This provides more proof that HSPD1 plays a crucial part in the pathogenetic processes that specifically cause stroke.

We used bioinformatics analysis to support the discovered link because no analysis of correlations of rs11682567 HSPD1 with the development of IS has been conducted so far. First, a large number of genes are impacted by the risk allele G of rs11682567 HSPD1's effects on their level of expression and alternative splicing, including: *ANKRD44*, which can regulate NF-κB, an important pro-inflammatory signaling pathway [24]; *PLCLI*, which modulates VEGFR2 signalling to promote angiogenesis in the central nervous system and vascular endothelial cell inflammation [25]; *MARS2*, regulating ROS production [26]; *RFTN2*, driving toll-like receptor 4 internalisation [27]; *COQ10B*, regulating mitochondrial respiratory chain assembly [28], brain ischemia-reperfusion [29] and ROS level [30]; *SF3B1*, a major regulator of hypoxia [31]; *HSPE1*, a component in ischemia damage control [32]. Second, it was discovered that the risk allele G of rs11682567 creates binding sites for transcription factors that jointly regulate biological processes with a high pathogenetic significance for IS. These biological processes encompass neurogenesis, monocyte differentiation, fatty acid metabolism, cell signaling (response to growth factor stimulus, to cytokine stimulus), and apoptosis. Third, it was discovered that the risk allele G is associated with an increase in diastolic blood pressure, total cholesterol levels, which are modifiable risk factors for the IS, as well as coronary heart disease, which is pathogenetically related to IS.

Numerous studies have shown that smoking can significantly modify the association of genetic markers with the risk of multifactorial diseases [33], including stroke [9-12]. The effect of cigarette smoke on the up-regulation of HSP60 in humans has been demonstrated in a large number of studies [34, 35]. This discovery further justifies the significance of endothelial stressors in initiating HSPD1-dependent pro-atherosclerotic molecular pathways and helps to explain the strong risk effects of rs11682567 *HSPD1* we observed in smokers and their absence in nonsmokers.

In conclusion, our case-control study findings indicate that the *HSPD1* gene variant of rs11682567 contributes to IS susceptibility. Further research is required to confirm these findings and look into the potential molecular mechanisms underlying the associations between *HSPD1* gene variants and IS risk.

Conclusion. The present study reveals the novel genetic association between rs11682567 *HSPD1* and the risk of IS observed exclusively in smokers. These data provide the new insights of involvement of HSP60 in cardiovascular pathology. Further studies may help to uncover the details of its particular role in the stroke pathogenesis.

Financial support

This research was funded by the Russian Science Foundation (№ 22-15-00288, <https://rscf.ru/en/project/22-15-00288/>).

Conflict of interests

The authors have no conflict of interest to declare.

References

1. GBD 2016 Neurology Collaborators. Global, regional, and national burden of neurological disorders, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet Neurology*. 2019;18(5):459-480. DOI: [https://doi.org/10.1016/S1474-4422\(18\)30499-X](https://doi.org/10.1016/S1474-4422(18)30499-X)
2. Bushueva O, Barysheva E, Markov A, et al. DNA Hypomethylation of the MPO Gene in Peripheral Blood Leukocytes Is Associated with Cerebral Stroke in the Acute Phase. *Journal of Molecular Neuroscience*. 2021;71(9):1914-1932. DOI: <https://doi.org/10.1007/s12031-021-01840-8>

3. Bushueva O, Solodilova M, Ivanov V, et al. Gender-specific protective effect of the -463G>A polymorphism of myeloperoxidase gene against the risk of essential hypertension in Russians. *Journal of the American Society of Hypertension*. 2015;9(11):902-906. DOI: <https://doi.org/10.1016/j.jash.2015.08.006>
4. Wick C. Heat Shock Protein 60: A Mediator of Atherosclerosis and Its Potential Therapeutic Role. In: Asea AAA, Kaur P, editors. *Chaperone Activity of Heat Shock Proteins*. Cham: Springer International Publishing; 2019. DOI: https://doi.org/10.1007/978-3-030-02254-9_4
5. Hu C, Yang J, Qi Z, et al. Heat shock proteins: Biological functions, pathological roles, and therapeutic opportunities. *MedComm*. 2022;3(3):e161. DOI: <https://doi.org/10.1002/mco2.161>
6. Kim JY, Kim JW, Yenari MA. Heat shock protein signaling in brain ischemia and injury. *Neuroscience Letters*. 2020;715:134642. DOI: <https://doi.org/10.1016/j.neulet.2019.134642>
7. Caruso Bavisotto C, Alberti G, Vitale AM, et al. Hsp60 Post-translational Modifications: Functional and Pathological Consequences. *Frontiers in Molecular Biosciences*. 2020;7:95. DOI: <https://doi.org/10.3389/fmolb.2020.00095>
8. Jakic B, Wick G, Cappellano G. Hsp60 in Atherosclerosis: Past, Present and Future. In: Asea AAA, Kaur P, editors. *Heat Shock Protein 60 in Human Diseases and Disorders*. Cham: Springer International Publishing; 2019. DOI: https://doi.org/10.1007/978-3-030-23154-5_13
9. Kobzeva KA, Shilenok IV, Belykh AE, et al. C9orf16 (BBLN) Gene, Encoding a Member of Hero Proteins, Is a Novel Marker in Ischemic Stroke Risk. *Research Results in Biomedicine*. 2022;8(3):278-292. DOI: <https://doi.org/10.18413/2658-6533-2022-8-3-0-2>
10. Belykh AE, Soldatov VO, Stetskaya TA, et al. Polymorphism of SERF2, the gene encoding a heat-resistant obscure (Hero) protein with chaperone activity, is a novel link in ischemic stroke. *IBRO Neuroscience Reports*. 2023;14:453-461. DOI: <https://doi.org/10.1016/j.ibneur.2023.05.004>
11. Shilenok I, Kobzeva K, Stetskaya T, et al. SERPINE1 mRNA Binding Protein 1 Is Associated with Ischemic Stroke Risk: A Comprehensive Molecular-Genetic and Bioinformatics Analysis of SERBP1 SNPs. *International Journal of Molecular Sciences*. 2023;24(10):8716. DOI: <https://doi.org/10.3390/ijms24108716>
12. Kobzeva KA, Soldatova MO, Stetskaya TA, et al. Association between HSPA8 Gene Variants and Ischemic Stroke: A Pilot Study Providing Additional Evidence for the Role of Heat Shock Proteins in Disease Pathogenesis. *Genes*. 2023;14(6):1171. DOI: <https://doi.org/10.3390/genes14061171>
13. Bushueva OY, Bulgakova IV, Ivanov VP, et al. Association of Flavin Monooxygenase Gene E158K Polymorphism with Chronic Heart Disease Risk. *Bulletin of Experimental Biology and Medicine*. 2015;159(6):776-778. DOI: <https://doi.org/10.1007/s10517-015-3073-8>
14. Bushueva OYu. Single Nucleotide Polymorphisms in Genes Encoding Xenobiotic Metabolizing Enzymes Are Associated with Predisposition to Arterial Hypertension. *Research Results in Biomedicine*. 2020;6(4):447-456. DOI: <https://doi.org/10.18413/2658-6533-2020-6-4-0-1>
15. Polonikov A, Bocharova I, Azarova I, et al. The Impact of Genetic Polymorphisms in Glutamate-Cysteine Ligase, a Key Enzyme of Glutathione Biosynthesis, on Ischemic Stroke Risk and Brain Infarct Size. *Life*. 2022;12(4):602. DOI: <https://doi.org/10.3390/life12040602>
16. Polonikov AV, Klyosova EYu, Azarova IE. Bioinformatic tools and internet resources for functional annotation of polymorphic loci detected by genome wide association studies of multifactorial diseases (review). *Research Results in Biomedicine*. 2021;7(1):15-31. Russian. DOI: <https://doi.org/10.18413/2658-6533-2020-7-1-0-2>
17. Wachoski-Dark E, Zhao T, Khan A, et al. Mitochondrial Protein Homeostasis and Cardiomyopathy. *International Journal of Molecular Sciences*. 2022;23(6):3353. DOI: <https://doi.org/10.3390/ijms23063353>
18. Duan Y, Tang H, Mitchell-Silbaugh K, et al. Heat Shock Protein 60 in Cardiovascular Physiology and Diseases. *Frontiers in Molecular Biosciences*. 2020;7:73. DOI: <https://doi.org/10.3389/fmolb.2020.00073>
19. Grundtman C, Kreutmayer SB, Almanzar G, et al. Heat shock protein 60 and immune inflammatory responses in atherosclerosis. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2011;31(5):960-968. DOI: <https://doi.org/10.1161/ATVBAHA.110.217877>
20. Saleki K, Alijanizade P, Moradi S, et al. Engineering a novel immunogenic chimera protein utilizing bacterial infections associated with atherosclerosis to induce a deviation in adaptive immune responses via Immunoinformatics approaches. *Infection, Genetics and Evolution*.

- 2022;102:105290. DOI: <https://doi.org/10.1016/j.meegid.2022.105290>
21. Rodriguez A, Von Salzen D, Holguin BA, et al. Complex Destabilization in the Mitochondrial Chaperonin Hsp60 Leads to Disease. *Frontiers in Molecular Biosciences*. 2020;7:159. DOI: <https://doi.org/10.3389/fmolb.2020.00159>
22. Vila-Casahonda RG, Lozano-Aponte J, Guerrero-Beltrán CE. HSP60-Derived Peptide as an LPS/TLR4 Modulator: An in silico Approach. *Frontiers in Cardiovascular Medicine*. 2022;9:731376. DOI: <https://doi.org/10.3389/fcvm.2022.731376>
23. Zhao Y, Zhang C, Wei X, et al. Heat shock protein 60 stimulates the migration of vascular smooth muscle cells via Toll-like receptor 4 and ERK MAPK activation. *Scientific Reports*. 2015;5:15352. DOI: <https://doi.org/10.1038/srep15352>
24. Poppenberg KE, Jiang K, Tso MK, et al. Epigenetic landscapes suggest that genetic risk for intracranial aneurysm operates on the endothelium. *BMC Medical Genomics*. 2019;12:149. DOI: <https://doi.org/10.1186/s12920-019-0591-7>
25. Kim S, Lee M, Choi YK. The Role of a Neurovascular Signaling Pathway Involving Hypoxia-Inducible Factor and Notch in the Function of the Central Nervous System. *Biomolecules and Therapeutics*. 2020;28(1):45-57. DOI: <https://doi.org/10.4062/biomolther.2019.119>
26. Mei X, Qi D, Zhang T, et al. Inhibiting MARSs reduces hyperhomocysteinemia-associated neural tube and congenital heart defects. *EMBO Molecular Medicine*. 2020;12(3):e9469. DOI: <https://doi.org/10.15252/emmm.201809469>
27. Yu H, Cheng W, Zhang X, et al. Integration analysis of methylation quantitative trait loci and GWAS identify three schizophrenia risk variants. *Neuropsychopharmacology*. 2020;45(7):1179-1187. DOI: <https://doi.org/10.1038/s41386-020-0605-3>
28. Guerra RM, Pagliarini DJ. Coenzyme Q biochemistry and biosynthesis. *Trends in Biochemical Sciences*. 2023;48(5):463-476. DOI: <https://doi.org/10.1016/j.tibs.2022.12.006>
29. Yuan B, Zhao XD, Shen JD, et al. Activation of SIRT1 Alleviates Ferroptosis in the Early Brain Injury after Subarachnoid Hemorrhage. *Oxidative Medicine and Cellular Longevity*. 2022;2022:9069825. DOI: <https://doi.org/10.1155/2022/9069825>
30. Wang Y, Hekimi S. The Complexity of Making Ubiquinone. *Trends in Endocrinology and Metabolism*. 2019;30(12):929-943. DOI: <https://doi.org/10.1016/j.tem.2019.08.009>
31. Hu Y, Lu H, Li H, et al. Molecular basis and clinical implications of HIFs in cardiovascular diseases. *Trends in Molecular Medicine*. 2022;28(11):916-938. DOI: <https://doi.org/10.1016/j.molmed.2022.09.004>
32. Liu F, Yang Y, Liu T, et al. Analysis of Differentially Expressed Long Noncoding RNA in Renal Ischemia-Reperfusion Injury. *Kidney and Blood Pressure Research*. 2020;45(5):686-701. DOI: <https://doi.org/10.1159/000508217>
33. Polonikov AV, Samgina TA, Nazarenko PM, et al. Alcohol Consumption and Cigarette Smoking are Important Modifiers of the Association Between Acute Pancreatitis and the PRSS1-PRSS2 Locus in Men. *Pancreas*. 2017;46(2):230-236. DOI: <https://doi.org/10.1097/MPA.0000000000000729>
34. Henderson B, Csordas A, Backovic A, et al. Cigarette smoke is an endothelial stressor and leads to cell cycle arrest. *Atherosclerosis*. 2008;201(2):298-305. DOI: <https://doi.org/10.1016/j.atherosclerosis.2008.02.022>
35. Kreutmayer SB, Messner B, Knoflach M, et al. Dynamics of heat shock protein 60 in endothelial cells exposed to cigarette smoke extract. *Journal of Molecular and Cellular Cardiology*. 2011;51(5):777-780. DOI: <https://doi.org/10.1016/j.yjmcc.2011.07.003>

Received 17 June 2023

Revised 2 September 2023

Accepted 27 November 2023

Information about the authors

Tatiana A. Stetskaya, Cand. Sci. (Biology), Junior Researcher at the Laboratory of Statistical Genetics and Bioinformatics, Research Institute for Genetic and Molecular Epidemiology, Kursk State Medical University, Kursk, Russia, E-mail: stetskaya_tatiana@mail.ru, ORCID: <https://orcid.org/0000-0002-4729-4654>.

Ksenia A. Kobzeva, Student, Kursk State Medical University, Kursk, Russia, E-mail: kseniya.kobzeva.0246@gmail.com, ORCID: <https://orcid.org/0000-0002-0652-2219>.

Sergey M. Zaytsev, Anesthesiologist-resuscitator, Pavlov First Saint Petersburg State Medical University, Saint Petersburg, Russia, E-mail: zukunftarzt.serii@yandex.ru, ORCID: <https://orcid.org/0000-0002-4280-9241>.

Irina V. Shilenok, Neurologist, Kursk City Clinical Emergency Hospital, Kursk, Russia, E-mail: irinka.zharikova.96@mail.ru, ORCID: <https://orcid.org/0000-0001-7894-8385>.

Galina V. Komkova, Cand. Sci. (Biology), Associate Professor at the Department of Biology, Medical Genetics and Ecology, Kursk State Medical University, Kursk, Russia, E-mail: komkovagv@kursksmu.net, ORCID: <https://orcid.org/0009-0003-2756-4164>.

Natalia V. Goryainova, Cand. Sci. (Biology), Associate Professor at the Department of Biology,

Medical Genetics and Ecology, Kursk State Medical University, Kursk, Russia, E-mail: goryainovanv@kursksmu.net, ORCID: <https://orcid.org/0009-0008-4980-4292>.

Olga Yu. Bushueva, Cand. Sci. (Medicine), Associate Professor at the Department of Biology, Medical Genetics and Ecology, Head of the Laboratory of Genomic Research, Research Institute for Genetic and Molecular Epidemiology, Kursk State Medical University, Kursk, Russia, E-mail: olga.bushueva@inbox.ru, ORCID: <https://orcid.org/0000-0003-3333-0623>.